

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/11, C07H 21/00, A61K 31/712 // A61P 35/00	A2	(11) International Publication Number: WO 00/24885 (43) International Publication Date: 4 May 2000 (04.05.00)
(21) International Application Number: PCT/US99/24758 (22) International Filing Date: 22 October 1999 (22.10.99) (30) Priority Data: 60/105,695 26 October 1998 (26.10.98) US (71) Applicant: AVI BIOPHARMA, INC. [US/US]; Suite 1105, One S.W. Columbia, Portland, OR 97258 (US). (72) Inventor: IVERSEN, Patrick, L.; 5902 N.W. Fair Oaks Place, Corvallis, OR 97330 (US). (74) Agents: GORTHEY, LeeAnn et al.; Dehlinger & Associates, P.O. Box 60850, Palo Alto, CA 94306-0850 (US).		(81) Designated States: AU, CA, JP, KR, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: p53 ANTISENSE AGENT AND METHOD (57) Abstract Antisense oligonucleotides useful for treating a disease state characterized by p53 induction, such as proliferative cell disorders, e.g. cancer, or a hypoxic state induced by an ischemic attack, such as stroke, are described. The antisense agents are preferably of the class known as "steric blocker" type oligonucleotides, including morpholino oligonucleotides, peptide nucleic acids, 2' O-allyl or 2'-O-alkyl modified oligonucleotides, or N3' → P5' phosphoramidate oligonucleotides.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

p53 ANTISENSE AGENT AND METHODField of the Invention

The present invention relates to antisense agents and methods for the treatment of disease states which are characterized by induction of p53. Such conditions include proliferative cell disorders, such as cancer, restenosis, and psoriasis, and hypoxic states induced by ischemic attacks such as stroke.

References

- Adhad, N. *et al.*, *Biochem. Biophys. Res. Commun.* 247:294-301 (1998).
- Agrawal, S. *et al.*, *Proc. Natl. Acad. Sci. USA* 87(4):1401-5 (1990).
- Arora, V., Ph.D. Dissertation, Dep. Pharmacol., Univ. Nebraska Coll. Med. (1998).
- Assy, N. *et al.*, "Use of proliferating cell nuclear antigen as a marker of liver regeneration after partial hepatectomy in rats," *J Lab Clin Med* 131(3):251-256 (1998).
- Bonham, M.A. *et al.*, *Nucleic Acids Res.* 23(7):1197-1203 (1995).
- Boudvillain, M. *et al.*, *Biochemistry* 36(10):2925-31 (1997).
- Chen, J.-K. *et al.*, *Nucleic Acids Res.* 23:2661-8 (1994).
- Cross, S.M. *et al.*, *Science* 267 (5202) p1353-6 (1995).I
- Edelman, E.R. *et al.*, *Circ. Res.* 76(2):176-182 (1995).
- Farrell, C.L. *et al.*, *Antisense Res. and Dev.* 5:175-183 (1995).
- Gambacorti-Passerini, C. *et al.*, *Blood* 88(4):1411-7 (1996).
- Gee, J.E. *et al.*, *Antisense & Nucleic Acid Drug Dev.* 8:103-111 (1998).
- Giles, R.V. *et al.*, *Anticancer Drug Des.* 8(1):33-51 (1993).
- Graeber, T.G. *et al.*, *Mol. Cell Biol.* 14(9):6264-77 (1994).
- Grisham, J.W. *et al.*, "Cell cycle and chemical carcinogenesis," *Surv. Synth. Pathol. Res.* 9-66 (1983).
- Gryaznov, S.M. *et al.*, *Proc. Natl. Acad. Sci. USA* 92:5798-5802 (1995).
- Hanvey, J.C. *et al.*, *Science* 258:1481-85 (1992).
- Hartwell, L.H. and Kastan, M.B. "Cell cycle control and cancer," *Science* 266:1821-28 (1994).
- Higgins, G.M. and Anderson, R.M. *Arch. Pathol.* 12:186. (1931).
- Horiguchi *et al.*, *Takeda Kenyushoho* 55:149-159 (1996).
- Johansson, H.E. *et al.*, *Nucleic Acids Res.* 22:4591-8 (1994).
- Kanoh *et al.*, *Bioorg. Chem. Lett.* 7:2847 (1997).
- Kastan, M.B. *et al.*, *Cell* 71: 587-97. (1992).
- Khooustov, V.I., *et al.*, *Lab. Invest.* 73:118-127 (1995).
- Knudsen, H. and Nielsen, P.E., *Nucleic Acids Res.* 24(3):494-500 (1996).

- Koch *et al.*, *Biochem. Biophys. Res. Commun.* 225:531-536 (1996).
- Levine, A.J. "p53, The cellular gatekeeper for growth and division," *Cell* 88:323-331 (1997).
- Li, Y. *et al.*, *Brain Res* 765 (2) p301-12 (1997).
- Lu, X. and Lane, D.P., *Cell* 75(4):765-78 (1993).
- 5 Magnelli, L. *et al.*, *Biochem Mol Med* 62(1):3-10 (1997).
- Messmer, U.K. *et al.*, "p53 Expression in nitric oxide-induced apoptosis", *FEBS Lett* 535(1):23-6 (1994).
- Nelson, W.G. and Kastan, M.B., *Mol. Cell Biol.* 14: 1815-1823 (1994).
- Nielsen, P.E. *et al.*, *Science* 254:1497-1503 (1991).
- 10 Raafat, A.M. *et al.*, *Shock Sep* 8 (3) p186-92 (1997).
- Rininger, J.A. *et al.* "Time course comparison of cell-cycle protein expression following partial hepatectomy and WY14,643-induced hepatic cell proliferation in F344 rats." *Carcinogenesis* 18(5):935-41 (1997).
- Rosenberg, R.D. *et al.*, U.S. Patent No. 5,593,974 (1997).
- 15 Russell, K.J. *et al.*, *Int. J. Radiat. Oncol. Biol. Phys.* 36(1099-1106 (1996).
- Russo, A. *et al.*, *Cancer Treatment Reports*, 69(11):1293-96 (1985).
- Sato, F. *et al.*, *Biochem. Biophys. Res. Commun.* 204:578-584 (1994).
- Sedlacek *et al.*, *Int. J. Oncol.* 9:1143-1168 (1996).
- Shu, M. *et al.*, *Apoptosis* 2:263-470 (1997).
- 20 Soini, T. *et al.*, *Br. J. Dermatol.* 131(4):514-20 (1994).
- Speir, E. *et al.*, *Scand. J. Infect. Dis. Suppl.* 99:78-81 (1995).
- Swe, M. *et al.*, *Biochem. Biophys. Res. Commun.* 236:594-598 (1997).
- Tadini, G. *et al.*, *Acta Derm. Venereol. Suppl.* 146:33-35 (1989).
- Van Dolah, F.M. *et al.*, *J. Cell Physiol.* 166:49-56 (1996).
- 25 Wagner, R.W. *et al.*, *Science* 260:1510-13 (1993).
- Waldman, T. *et al.*, "Uncoupling of S phase and mitosis induced by anticancer agents in cells lacking p21," *Nature* 381:713-716 (1996).
- Xiang, H. *et al.*, *J Neurosci* 16 (21) p6753-65 (1996).
- Zhang, M. and Raveche, E.S., *Oncol Rep* 5 (1) p23-30 (1998).
- 30 Zhu, W.Y. *et al.*, *Exp. Cell Res.* 2:463-470 (1997).

Background of the Invention

The importance of p53 in cell proliferation is apparent from the observation that more than half of all human cancers display p53 mutations (Levine, 1997). An induction in p53 expression is observed in cells exposed to DNA alkylating agents, nitric oxide (Messmer, 1994), ionizing and ultraviolet irradiation (Lu and Lane, 1993), the restriction enzyme PvuII, which induces

chromosome damage (Lu and Lane, 1993), and other agents capable of inducing DNA strand breaks (Nelson and Kastan, 1994). Ischemia/reperfusion (Raafat, 1997), models of epilepsy (Xiang, 1996), and hypoxia (Graeber, 1994) have also been demonstrated to induce p53. As shown herein and in previous studies (Rininger, 1997), p53 protein levels are also highly upregulated during the course of liver regeneration following partial hepatectomy, showing that damage of endogenous origin can also cause induction of p53 *in vivo*.

Roles for p53 have been postulated in cell cycle checkpoint activity, apoptosis, differentiation and DNA repair (Magnelli, 1997). Cell cycle checkpoints serve two important functions. One is to ensure that essential events in the cell cycle are completed before a subsequent event; the other is to provide more time for repair of damaged DNA before DNA replication and mitosis occur (Hartwell, 1994). The checkpoint activity of p53 occurs predominantly at the level of cells entering the S-phase (DNA synthesis) of the cell cycle (Kastan, 1992). G₂-M checkpoint activity has also been reported for p53, due to its role as an upstream regulator of p21^{waf-1}. In the absence of either protein, DNA-damaged cells arrest in a G₂-like state, but then undergo additional S phases without intervening normal mitoses (M phase), leading to nuclear abnormalities, e.g. polyploidy, and culminating in apoptosis (Waldman, 1996).

Summary of the Invention

The invention provides, in one aspect, methods and compositions for treating a disease state characterized by p53 induction in a mammalian subject. The method comprises administering to the subject, in a suitable pharmaceutical carrier, a pharmaceutically effective amount of an antisense agent having the base sequence SEQ ID NO:1, identified as 5'-TCA GTC TGA GTC AGG CCC-3', or the base sequence SEQ ID NO:2, identified as 5'-CCC TGC TCC CCC CTG GCT CC-3', where the antisense agent is a morpholino oligonucleotide, a peptide nucleic acid, a 2'-O-allyl or 2'-O-alkyl modified oligonucleotide, a N3'→P5' phosphoramidate oligonucleotide, or a C-5-propyne pyrimidine-modified oligonucleotide. In a preferred embodiment, the antisense agent is a morpholino oligonucleotide, preferably having morpholino subunits joined by phosphorodiamidate backbone linkages. In another preferred embodiment, the antisense agent is a C-5-propyne pyrimidine-modified oligonucleotide.

The invention also includes compositions for use in treating such disease states. These compositions include antisense agents as described above, in a suitable pharmaceutical carrier.

In preferred applications of the method, the subject is a human subject. The disease state may be one which results from an ischemic or ischemic/reperfusion injury, such as stroke or the aftermath of organ transplant. Alternatively, the disease state may be cancer, in which case a preferred embodiment further comprises administering an agent effective to increase radical oxygen species at the cellular level. Such agents include radiosensitizing agents, ionizing radiation, a

hyperbaric oxygen environment, and chemotherapeutic agents, such as certain anthracyclines or anthraquinones, which increase radical oxygen species at the cellular level.

For cancer treatment, the method may also include administering an agent effective to interfere with progression from the G₂ phase to the M phase of the cell cycle. Such agents include phosphokinase C (PKC) inhibitors, bis(chloroethyl)nitrosourea (BCNU), pentoxifylline, silymarin, staurosporine, phenylahistin, paclitaxel, retinoic acid, flavopiridol, methyl-2,5-dihydrocinnamate, herboxidiene, 9-nitrocamptothecin, maitotoxin, apigenin, nocodazole, and colcemid.

In another aspect, the invention provides an oligonucleotide having a base sequence selected from the group consisting of SEQ ID NO:1, identified as 5'-TCA GTC TGA GTC AGG CCC-3', and SEQ ID NO:2, identified as 5'-CCC TGC TCC CCC CTG GCT CC-3', where the oligonucleotide is selected from the group consisting of a morpholino oligonucleotide, a peptide nucleic acid, a 2'-O-allyl or 2'-O-alkyl modified oligonucleotide, and a N3' → P5' phosphoramidate oligonucleotide. In one embodiment, the oligonucleotide is a morpholino oligonucleotide, preferably comprising morpholino subunits joined by phosphorodiamidate backbone linkages. The invention also provides an oligonucleotide having a base sequence selected from the group consisting of SEQ ID NO:1, identified as 5'-TCA GTC TGA GTC AGG CCC-3', and SEQ ID NO:2, identified as 5'-CCC TGC TCC CCC CTG GCT CC-3', where the oligonucleotide is a C-5-propyne pyrimidine-modified oligonucleotide.

These and other objects and features of the invention will become more fully apparent when the following detailed description of the invention is read in conjunction with the accompanying drawings.

Brief Description of the Drawings

Fig. 1 shows the structure of a phosphorodiamidate-linked morpholino oligonucleotide analog; and

Fig. 2 shows the wet weights (in gm) of remnant regenerating livers in rats, 24 hr after partial hepatectomy and administration of the following antisense oligonucleotides, having SEQ ID NOs 1 and 2, respectively: a phosphorothioate (S-ODN, 1 mg/200g); a C-5-P modified phosphorothioate (C-5-P, 0.1mg/200g); and a neutral backbone (phosphorodiamidate) morpholino oligonucleotide (3.4 mg/200g). Error bars represent statistical difference of $p < 0.05$ with respect to the saline-treated control of each respective group.

Detailed Description of the Invention

I. Definitions

An "antisense oligonucleotide" or "antisense agent" refers to a molecule which includes a sequence of purine and pyrimidine heterocyclic bases, supported by a backbone, which are

effective to hydrogen-bond to corresponding, contiguous bases in a target nucleic acid sequence. The backbone is composed of subunit backbone moieties supporting the purine and pyrimidine heterocyclic bases at positions which allow such hydrogen bonding. These backbone moieties are cyclic moieties of 5 to 7 atoms in length, linked together by phosphorous-containing linkages one to three atoms long.

A "morpholino" oligonucleotide is composed of morpholino subunit structures of the form shown in Fig. 1, where (i) the structures are linked together by phosphorous-containing linkages, one to three atoms long, joining the morpholino nitrogen of one subunit to the 5' exocyclic carbon of an adjacent subunit, and (ii) B is a purine or pyrimidine base-pairing moiety effective to bind, by base-specific hydrogen bonding, to a base in a polynucleotide. Fig. 1 illustrates two such subunits joined by a phosphorodiamidate linkage.

A "N3' → P5' phosphoramidate" oligonucleotide is one in which the 3'-oxygen of the 2'-deoxyribose is replaced by a 3'-amine, as described, for example, by Gryaznov *et al.* and Chen *et al.*

A "2'-O-allyl (or alkyl) modified oligonucleotide" is an oligoribonucleotide in which the 2' hydroxyl is converted to an allyl or alkyl ether. The alkyl ether is typically a methyl ether.

A "C-5-propyne pyrimidine-modified oligonucleotide" is an oligonucleotide in which the C-5 methyl group of thymidine bases and/or the C-5 hydrogen of cytidine bases has been replaced with a propyne group.

In a "peptide nucleic acid", the deoxyribose phosphate units of an oligonucleotide backbone are replaced with polyamide linkages. As described, for example, in Nielsen *et al.* and Hanvey *et al.*, proper backbone spacing is attained by the use of 2-aminoethyl glycine units, with a nucleotide base attached to each 2-amino group via a methylenecarbonyl group.

A "RNase-inactive" oligonucleotide or oligonucleotide analog is one which acts via an RNase-independent mechanism, unlike RNase-active oligonucleotides, such as phosphorothioates. They are believed to function by sterically blocking target RNA formation, nucleocytoplasmic transport or translation, and are thus also referred to as "steric blockers". This class includes, for example, methylphosphonates, morpholino oligonucleotides, as described herein, peptide nucleic acids (PNA's), 2'-O-allyl or 2'-O-alkyl modified oligonucleotides, and N3' → P5' phosphoramidates.

A condition "characterized by induction of p53" is a disease state in which levels or expression of p53 protein in affected cells are increased relative to the non-disease state, and/or a condition in which suppression of p53 expression, sometimes in combination with other treatment, has a beneficial therapeutic effect. Examples are cancer, ischemia, such as results from stroke, myocardial infarction, or epileptic seizure, and the resulting hypoxic state, and ischemic/reperfusion injury, such as may occur in organ transplants.

Abbreviations:

ON = oligonucleotide

ODN = oligodeoxyribonucleotide

S-ODN = phosphorothioate oligonucleotide

5 C-5-P = C-5-propyne pyrimidine-modified oligonucleotide

II. Antisense OligonucleotidesA. Sequences

10 The antisense agents of the present invention comprise nucleotide subunits joined by internucleotide backbone linkages which present the nucleotide bases for hybridization with target RNA sequences. The base sequences of these agents are complementary (antisense) to portions of p53 mRNA. The first sequence, corresponding to SEQ ID NO: 1, is complementary to a unique portion of rat p53 mRNA (residues 1182 to 1199; Genbank accession # X13058) and is referred to herein as "p53T". This sequence is the same in rat, mouse, monkey, and human
15 mRNA, and thus ideally suited for *in vivo* testing in animal models.

The second sequence, named "OL(1)p53" and corresponding to SEQ ID NO: 2, is antisense to human p53, and is therefore effective at inhibiting p53 expression in humans (Iverson, U.S. Patent No. 5,641,754). Because it has four mismatches with the rat sequence, it does not alter such expression in rats, and therefore provides a useful control in rat model studies.

20

B. Oligonucleotide Structure

Various oligonucleotide analogs known in the art present advantages over "natural" polynucleotides in areas such as stability, particularly nuclease resistance, reduction of non-specific binding, and bioavailability (i.e. access to cells). The structures may be modified at the
25 backbone, the sugar moiety, or the bases themselves. Such analogs include, for example, phosphorothioates (designated herein as S-ODN), methylphosphonates, phosphotriesters, C-5-propyne pyrimidine-modified oligonucleotides (C-5-P), morpholino oligonucleotides, peptide nucleic acids (PNA's), 2'-O-allyl or 2'-O-alkyl modified oligonucleotides, and N3' → P5' phosphoramidates.

30 Preparation of such antisense agents is well known in the art, and may often be conveniently carried out on automated synthesizers. A general procedure for synthesis of S-ODN, C-5-P, and uncharged morpholino antisense oligonucleotides is given in Example 1.

Two general mechanisms have been proposed to account for inhibition of expression by antisense oligonucleotides (see *e.g.* Agrawal; Bonham; Boudvillain; and references cited therein).
35 In the first, a heteroduplex formed between the oligonucleotide and mRNA serves as a substrate for RNase H, leading to cleavage of the mRNA. Oligonucleotides belonging, or proposed to

belong, to this class include phosphorothioates, phosphotriesters, and phosphodiester (i.e., unmodified "natural" oligonucleotides). Such compounds generally show high activity, and phosphorothioates are currently the most widely employed oligonucleotides in antisense applications. However, these compounds tend to produce unwanted side effects due to non-specific binding to cellular proteins (Gee), as well as inappropriate RNase cleavage of non-target RNA heteroduplexes (Giles).

A second class of oligonucleotide analogs, termed "steric blockers" or, alternatively, "RNase inactive" or "RNase resistant", are not observed to activate RNase H, and are believed to act by sterically blocking target RNA formation, nucleocytoplasmic transport or translation. This class includes methylphosphonates (Toulme) as well as morpholino oligonucleotides, peptide nucleic acids (PNA's), 2'-O-allyl or 2'-O-alkyl modified oligonucleotides (Bonham), and N3' → P5' phosphoramidates (Gee). They are reported to be most effective when targeted at the AUG start codon, the 5'-splice site, or the 5'-untranslated region of mRNA's. To date, they have shown relatively weak (Bonham) or, more often, no activity when targeted downstream, at coding regions (see, for example, Toulme, for methylphosphonates; Gambacorti, for PNA's; Knudsen, for duplex-forming PNA's; and Gee, for PNA's and N3' → P5' phosphoramidates). It has been proposed that the hybrids formed by these ON's, when directed to the coding region, are not sufficiently stable to avoid being displaced by the unwinding action of the ribosomal complex during translation (Boudvillain, Gee). One approach to addressing this problem has been to covalently link or intercalate the ON to its target sequence (Gee, Johansson). However, without such modifications, the activity of steric blocking ON's has generally been limited to the initiation regions noted above.

III. Induction of p53 in Proliferating Cells and Suppression by Antisense p53

In previously reported studies (Arora, 1998), antisense agents p53T and ON(1)p53, described above, were administered to rats immediately after partial hepatectomy surgery, as described in Example 2. The regenerating rat liver is an excellent *in vivo* model system to study the mechanisms of growth control and cellular proliferation in a natural tissue environment. The rapidly proliferating cells in the regenerating liver are extremely susceptible to carcinogenesis, as there is minimal time available between cell cycles for DNA repair (Grisham, 1983). As noted above, ON(1)p53 does not alter p53 expression in rats, and was used as a control.

A massive induction of p53 was observed during the process of regeneration in the PH (post-hepatectomy) rats. The following effects of antisense ON administration were measured: weight gain of regenerating liver; level of p53, PCNA, p21 and NADPH; DNA content of isolated cells; microsomal protein content; various microsomal enzyme assays; oxidative stress, as evidenced by lipid peroxidation; and mitotic index.

When treated with antisense p53T (SEQ ID NO:1), the livers lost their G₁-S cell cycle checkpoint activity, as evidenced by increases (relative to controls) in weight gain, mitotic index, and PCNA expression, which is an accurate marker of cell proliferation (Assy, 1998), and a 5-fold reduction in G₁ cell population. The p53T-treated PH rat livers also displayed a greater number of multi-nucleate cells. Functional recovery of the regenerating liver, as determined by enzyme assays, was also increased by p53 inhibition.

In summary, the data showed that p53 expression is inhibited in regenerating livers of post-hepatectomy rats treated with antisense p53T (SEQ ID NO: 1), and that this inhibition results in enhanced mitosis, elevated PCNA expression, and diminished number of cells in the G₁ phase of the cell cycle. The results are consistent with G₁-S cell cycle checkpoint activity of p53.

IV. Comparative Effect of S-ODN, C-5-P, and Morpholino Antisense Agents

Neutral-backbone morpholino antisense agents having the base sequences given as SEQ ID NO: 1 and SEQ ID NO: 2 were prepared as described in Example 1. Such morpholino oligomers have shown high binding affinity for RNA targets, and the uncharged backbone favors uptake into cells and reduces non-specific binding interactions, relative to charged analogs such as phosphorothioates. Also prepared were the corresponding C-5-propyne cytosine modified (C-5-P) phosphorothioates. As discussed above, the morpholino analog is believed to act as a "steric blocker". C-5-P phosphorothioates are reported to act by an RNase-H dependent mechanism (Wagner), attributed to phosphorothioates in general.

The effect of intraperitoneal administration of these antisense agents was compared to that of the unmodified phosphorothioates (S-ODNs) with respect to wet weight gain of remnant regenerating livers 24 hr post PH, as described in Section III and Example 2. The data, shown in Fig. 2, indicate that the p53T sequence is active in the liver regeneration model for both of the alternate analogs tested, and the OL(1)p53 sequence remained inactive (as expected) with the alternate analogs at the doses tested.

Efficacy similar to a 1 mg/200 gm dose of S-ODN p53T was observed with 100 µg/200 gm of the C-5-P modified S-ODN and 50 nM/200 gm (approximately 3.7 mg/200g) of the morpholino anti-p53T oligonucleotide (having SEQ ID NO: 1). It is especially noteworthy that the morpholino antisense oligomer is active when targeted to exon 10 of the coding region of the gene, rather than to the 5'-untranslated region or the AUG start codon. As noted above, "steric blocker" antisense agents generally have not shown significant activity when targeted downstream from the AUG translation initiation site. This coding sequence may be particularly favorable for antisense ON binding, forming duplexes stable enough to sterically block translation. In addition, as noted above, morpholino oligomers have proven to be particularly effective RNA-binding molecules. While the invention is not limited to a particular mechanism, it is postulated

that the RNA helix is distorted by the binding of the morpholino oligonucleotide, and that this distortion interferes with ribosomal binding.

Studies in support of the invention have suggested that the propyne groups of the C-5-P propyne modified ODN protrude into the major groove of the RNA duplex, and thus may also sterically interfere with ribosomal binding. Accordingly, substitution of other sterically bulky groups for the propyne group, including groups having different electronic structures, such as t-butyl, could produce a similar inhibitory effect.

V. Therapeutic Methods

The present invention provides a method for inhibiting p53 expression, particularly in cells characterized by p53 induction, and where inhibition of p53 produces a therapeutic benefit. The method can be used for treating a disease state characterized by p53 induction. The method comprises administering to the subject having such a condition, or to cells harvested from such a subject, an oligonucleotide with a sequence corresponding to SEQ ID NO: 1 or SEQ ID NO: 2. Preferably, the subject is a human subject. (As noted above, SEQ ID NO: 2 is not effective in rat models.) The oligonucleotide is preferably a morpholino oligonucleotide, a peptide nucleic acid, a 2'-O-allyl or 2'-O-alkyl modified oligonucleotide, a N3' → P5' phosphoramidate oligonucleotide, or a C-5-propyne pyrimidine-modified phosphorothioate oligonucleotide. Morpholino oligonucleotides are particularly preferred for both sequences. Also contemplated are C-5-P modified morpholino oligonucleotides.

As demonstrated by the data shown herein and reported in Arora, 1998, the anti-p53 ON suppress the expression of p53 in post-hepatectomy rats. Morpholino and C-5-P ON's were found to be more effective than equivalent amounts of the unmodified phosphorothioate (S-ODN), as determined by weight gain of the regenerating liver. The morpholino oligomer is surprisingly effective in view of the fact that it is targeted at a coding region (exon 10), well downstream of the AUG start region, the latter of which is the conventional target for RNase-inactive ("steric blocker") antisense oligonucleotides.

A. Anti-proliferative Therapy

As noted above, an induction in p53 expression is observed in cells exposed to various DNA-damaging agents. The damaged DNA is most commonly repaired by unscheduled DNA synthesis. Such repair normally occurs during the G₁ cell cycle phase, which precedes the S (synthesis) phase. If, however, the modified DNA bases remain unrepaired (e.g. by eliminating the G₁-S cell cycle checkpoint, thus sending the cell prematurely into the S phase), consequences can include mutagenesis, due to erroneous replication of damaged DNA nucleotide base templates, and/or cell death, due to inability of the cell to replicate its genome past a damaged

DNA site.

Inhibiting the expression of p53 suppresses this cell cycle checkpoint and can thus be used to selectively kill DNA-damaged cells. It was shown in U.S. Patent No. 5,641,764 (Iverson) that administration of an antisense phosphorothioate oligonucleotide having the sequence OL(1)p53
5 (SEQ ID NO: 2) produced apoptosis *ex vivo* and *in vivo* in human liver cells. The effect was increased by exposure of the cells to conditions which increased the net reactive oxygen content in the cells. Such treatment produces little or no adverse effect on normal tissue, as normal tissue has greater oxygen scavenging activity than cancer cells. Most cells, for example, normally contain one or more enzymes, such as SOD, catalase, or glutathione peroxidase, which
10 very rapidly combine with and inactivate excess reactive oxygen species. As described above, it is believed that the suppression of the G₁-S checkpoint by inhibition of p53 prevents repair of damaged DNA, leading to mutagenesis and cell death.

In a preferred method, therefore, particularly useful against cancer cells, the ON is administered in combination with an agent capable of increasing radical oxygen induced
15 cytotoxicity. The oligonucleotide and agent can be administered substantially concomitantly, or sequentially, with either therapeutic agent being given first. However, best results are achieved when the oligonucleotide is administered sufficiently in advance to permit therapeutic blood levels to be achieved.

Increased radical oxygen levels are produced, for example, by exposure of the cell to
20 radiation, or an agent capable of radical oxygen induced cytotoxicity, such as for example, anthracycline cytotoxic antibiotics (e.g., doxorubicin), BCNU, BSO (buthionine sulfoxamine), hydrogen peroxide, or antisense oligonucleotide inhibitors of SOD (superoxide dismutase), catalase, GSH synthetase, GSH reductase, or GSH peroxidase.

The agent capable of radical oxygen induced cytotoxicity can be, for example, a
25 radiosensitizer, a chemotherapeutic agent which generates radical oxygen, or an oligonucleotide capable of interaction with cells of the host causing the formation of hydroxy radicals (e.g., see Iversen, U.S. Ser. No. 07/735,067, filed Jul. 25, 1991, entitled "Inhibition of Mutagenicity Induced by Binding of Oligonucleotides to Cells", the content of which is incorporated herein by reference). A number of cytotoxic agents that have a lethal effect on target cells within the
30 patient, wherein the lethal effect may be enhanced by one or more sensitizing agents, may be used in the method of the invention. Included are radiation (whether the radiation therapy is delivered internally or administered by external means) and cytotoxic drugs. Examples of some of the many such drugs are nitrogen mustards such as L-phenylalanine nitrogen mustard (melphalan), anthracycline chemotherapeutics such as daunorubicin and doxorubicin, and
35 platinum compounds such as cis-diaminodichloro platinum (cis-platin).

Internally delivered radiation includes therapeutically effective radioisotopes injected into a patient. Such radioisotopes include, but are not limited to, the radionuclide metals ^{186}Re , ^{188}Re , ^{64}Cu , ^{67}Cu , ^{109}Pd , ^{212}Bi , ^{203}Pb , ^{212}Pb , ^{211}At , ^{97}Ru , ^{105}Rh , ^{198}Au , ^{199}Ag , and ^{131}I . These radioisotopes generally will be bound to carrier molecules (e.g., in the form of a chelate-antibody conjugate) when administered to a patient. Examples of suitable internally delivered radiotherapeutic agents are metal radionuclide chelates conjugated to antibodies as described, for example, in EP Publication No. 188,256. Radiation administered by external means includes external beam radiation such as cobalt therapy.

The choice of sensitizing agent depends on such factors as the particular type of tumor to be treated and the cytotoxic agent to be administered. Preferably, the compound is selective for a certain target site *in vivo*, such as a cancer cell. For example, U.S. Pat. No. 4,628,047 reports the use of diltiazem to enhance the sensitivity of a variety of types of cancer cells toward cytotoxic agents such as doxorubicin. Other pairings of sensitizers with cytotoxic drugs, as well as expected differences in the susceptibility of different types of cancer cells to treatment with these agents, are discussed in *Important Advances in Oncology* (DeVita *et al.*, editors, J.B. Lippincott Co., Philadelphia (1986), pp. 145-157).

A preferred sensitizer is BSO (buthionine sulfoxamine, available from Chemical Dynamics Corporation, South Plainfield, N.J.), a synthetic amino acid that inhibits γ -glutamylcysteine synthetase and leads to a marked decrease of glutathione (GSH) in cells. Thus, BSO is thought to function as a sensitizer for drugs that have a cytotoxic effect enhanced by decreasing cellular glutathione levels. Especially preferred is the S-isomer of BSO, which has utility in depleting glutathione in the cell (see U.S. Pat. Nos. 5,171,885 and 5,245,077).

Additional combinations of sensitizers and cytotoxic agents may be identified through such methods as *in vitro* assays using cultured cells which correspond to the desired target cells (e.g., a specific cancer cell line). For example, Russo *et al.* describe assays for determining whether BSO is effective in lowering glutathione synthetase levels in a particular type of cell line.

The above discussion of radical oxygen generating species, cytotoxic agents and sensitizers is primarily directed to the preferred target cells, i.e., cancer cells characterized by p53 expression. Such cancers include those of the bladder, brain, breast, cervix, colon, esophagus, larynx, liver, lung, ovary, pancreas, prostate, skin, stomach and thyroid.

In another preferred method, administration of the antisense ON is accompanied by administration of an agent effective to interfere with progression from the G_2 phase to the M (mitosis) phase of the cell cycle. Such agents include, for example, pentoxifylline (Russell, 1996), silymarin (Ahmad, 1998), staurosporine (Swe, 1997), phenylahistin (Kanoh, 1997), paclitaxel (Shu, 1997), retinoic acid (Zhu, 1997), flavopiridol (Sedlacek, 1996), methyl-2,5-dihydrocinnamate (Koch, 1996), herboxidiene (Horiguchi, 1996), 9-nitrocamptothecin (Khooustov, 1995), maitotoxin

(VanDolah, 1996), apigenin (Sato, 1994), nocodazole, and colcemid (Zhang, 1998). Recent reports (Cross, 1995) have suggested that p53 is instrumental in a spindle checkpoint, which ensures complete assembly of the spindle apparatus during the mitosis (M) phase of the cell cycle. In studies in support of the invention, p21-deficient mice, when treated with G₂-M inhibitors such as those listed above, showed a high degree of aneuploidy in the affected cells. It is postulated that the inhibitors prevented the cells from undergoing normal progression through the mitosis phase, and, lacking the mitotic (and G₁-S phase) checkpoint, the cells underwent successive synthesis (S) phases, resulting in aneuploidy. Because p53 is necessary for transcriptional activation of p21, cells in which p53 has been inhibited by the antisense agents of the invention should show a similar effect.

10 Spindle inhibitors such as nocodazole or colcemid (Cross, Zhang) may also be used in conjunction with anti-p53 antisense agents. In the absence of the p53-dependent spindle checkpoint, such cells are expected to undergo successive cell cycles with abnormal or incomplete mitotic phases, also leading to polyploidy or aneuploidy.

Other hyperproliferative disorders may also be treated. Examples of hyperproliferative skin disorders include human papilloma virus (HPV)-infected cells, commonly associated with warts, superficial neoplasias of the skin such as melanomas, pre-malignant and malignant carcinomas, actinic keratosis, lupus erythematosus, dermatitis, and psoriasis. Hyperproliferative diseases of other epithelial tissues (e.g. endothelium, mesothelium) include reversible obstructive airway disease, such as asthma and bronchitis. Various hyperproliferative disorders of the eye, including conical cornea, keratoconjunctivitis, vernal conjunctivitis, and keratitis leukoma, may also be treated. The methods may also be used to treat hyperproliferative vascular diseases such as intimal smooth muscle cell hyperplasia, vascular occlusion, and restenosis.

B. Treatment of Hypoxic or Ischemic States

25 Other disease states characterized by p53 induction include hypoxic states such as follow an ischemic attack such as a stroke. For example, p53 was observed to be preferentially expressed in apoptotic cells after ischemia in the cerebrum (Li), and in damaged neurons in models of ischemia and epilepsy (Xiang). Raafat observed increased levels of p53 following ischemia/reperfusion in kidneys, and proposed that the increase may facilitate apoptosis. Heat shock has also been reported to induce p53 (Graeber). Therefore, in another aspect, the invention includes treatment of tissues subjected to such stresses, e.g., heat shock, epileptic seizure, stroke, and other occurrences of ischemia and/or reperfusion (e.g. following organ transplant) with anti-p53 antisense agents as described herein.

VI. Formulations and AdministrationA. Therapeutic Formulations

For administration in accordance with the treatment methods of the invention, the antisense oligonucleotides are preferably combined with a pharmaceutically acceptable carrier, such as a
5 suitable liquid vehicle or excipient, and optional auxiliary additives. The liquid vehicles and excipients are conventional and commercially available. Illustrative thereof are distilled water, physiological saline, aqueous solutions of dextrose, and the like.

In general, in addition to the active compounds, the pharmaceutical compositions of the invention may contain suitable excipients and auxiliaries which facilitate processing of the active
10 compounds into preparations which can be used pharmaceutically. Suitable excipients are, in particular, fillers such as sugars, for example, lactose or sucrose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, and binders such as starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone. If desired, disintegrating agents may be added, such as the above-
15 mentioned starches as well as carboxymethyl starch, cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof. Auxiliaries include flow-regulating agents and lubricants, for example, silica, talc, stearic acid or salts thereof, and/or polyethylene glycol.

Dragee cores may be provided with suitable coatings which, if desired, may be resistant to gastric juices. For this purpose, concentrated sugar solutions may be used, which may optionally
20 contain gum arabic, talc, polyvinylpyrrolidone, polyethyleneglycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethyl-cellulose phthalate, are used. Dyestuffs and pigments may be added to the tablets of dragee coatings for identification or in order to characterize different
25 combinations of active compound doses.

Other pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer such as glycerol or sorbitol. The push-fit capsules can contain the active compounds in the form of granules which
30 may be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added.

Pharmaceutical preparations which can be used rectally include, for example, suppositories, which consist of a combination of the active compounds with a suppository base. Suitable
35 suppository bases include natural or synthetic triglycerides, paraffin hydrocarbons, polyethylene glycols, or higher alkanols. In addition, it is possible to use gelatin rectal capsules which consist

of a combination of the active compounds with a base. Possible base materials include liquid triglycerides, polyethylene glycols, or paraffin hydrocarbons.

Suitable liquid formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble or water-dispersible form. In addition, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, and/or dextran. The suspension may also contain stabilizers.

In addition to administration with conventional carriers, the active ingredients may be administered by a variety of specialized delivery techniques. For example, the compounds of the present invention may be administered encapsulated in liposomes. The active ingredient, depending upon its solubility, may be present both in the aqueous phase and in the lipidic layer(s), or in what is generally termed a liposomic suspension. The lipidic layer generally comprises phospholipids, such as lecithin or sphingomyelin, steroids such as cholesterol, ionic surfactants such as diacetylphosphate, stearylamine, or phosphatidic acid, and/or other hydrophobic materials. The diameters of the liposomes generally range from about 15 nm to about 5 microns.

Methods for preparing such dosage forms are known or will be apparent to those skilled in the art; for example, see Remington's Pharmaceutical Sciences (19th Ed., Williams & Wilkins, 1995). The pharmaceutical preparations are manufactured according to procedures well known in the art. For example, they may be made by means of conventional mixing, granulating, dragee-making, dissolving, or lyophilizing processes. The process to be used will depend ultimately on the physical properties of the active ingredient used.

B. Treatment of Tumors

The composition may be administered to a subject orally, transdermally or parenterally, e.g., by intravenous, subcutaneous, intraperitoneal, or intramuscular injection. For in vivo antineoplastic use, the p53 mRNA antisense oligonucleotides are preferably administered intravenously.

In accordance with the present invention, the oligonucleotide which inhibits expression of p53, thus acting as an oxidative repair path down-regulator, is administered to a patient prior to or concomitant with administration of the agent capable of radical oxygen induced cytotoxicity. When administered simultaneously, the two active agents can be in conjugated or unconjugated form. If in conjugated form, it is preferred that the conjugate comprises a cleavable linkage so that the oligonucleotide and the cytotoxic agent are released at the target site.

The amount of each agent administered is such that the combination of the two types of agents is therapeutically effective. Dosages will vary in accordance with such factors as the age, health, sex, size and weight of the patient, the route of administration, the toxicity of the drugs, and the relative susceptibilities of the cancer to the oligonucleotide and cytotoxic agent.

5 Recommended dosages and dosage forms for most of the cytotoxic agents have been established and can be obtained from conventional sources, such as the *Physicians Desk Reference*, published by Medical Economics Company, Inc., Oradell, N.J. If necessary, these parameters can be determined for each system by well-established procedures and analysis, e.g. in clinical trials.

For in vivo use, the preferred dosage of the oligonucleotides of the invention is typically that
10 which is necessary to attain a concentration in the blood of from about 0.01 to about 1 $\mu\text{mol/l}$. This concentration can be achieved in a variety of ways; doses of between about 0.05 and about 0.2mg/kg/hour by continuous IV infusion have been found to be acceptable. Greater or lesser amounts of oligonucleotide may be administered as required.

Example 3, below, demonstrates the application of the invention in the treatment of human
15 B-cell lymphoma.

C. Treatment of Ischemic Injury

The ischemic condition may be due to an interruption in cerebral circulation, such as caused by cardiac failure, or other condition leading to global loss of blood supply to the brain, or to localized
20 interruptions in blood flow, such as due to cerebral hemorrhaging, or localized thrombotic or embolic events, or head trauma. The ischemic condition to be treated is generally associated with stroke, defined as the sudden diminution or loss of neurological function caused by an obstruction or rupture of blood vessels in the brain. Secondary brain damage resulting from the original ischemic event typically includes cerebral cell destruction, or lesions, in the area surrounding the ischemic
25 injury, in the case of focal ischemia, and also in areas of selective vulnerability in lesions, such as the hippocampus or basal ganglia, in the case of global ischemia. The secondary damage may often be manifested by functional impairment, such as loss of short-term or long-term memory.

The ON is formulated for parenteral administration in a suitable inert carrier, such as a sterile physiological saline solution. The dose administered will be determined by route of administration.
30 One suitable route is intracerebroventricular (ICV), at a dose level of about 50 μg - 5 mg ON/kg body weight. A pharmaceutically effective dose, i.e., a dose effective to produce significant reduction in anatomical and/or functional damage, can be estimated from the dose/response seen in model systems for ischemia and stroke damage. Such model systems include the gerbil model of global ischemia, produced by transient occlusion of carotid arteries of the neck, and the rat four-
35 vessel occlusion model. Tissues are examined for anatomical damage, and functional damage may

be assessed by observation of hyperactivity, a common consequence of cerebral ischemia in animals, or effect on short-term memory. These procedures are known in the art ; see for example, U.S. Patent No. 5,051,403.

5 D. Treatment of Restenosis

 An important aspect of successful anti-restenosis therapy, particularly with regard to antisense agents, is effective delivery of the antisense oligomer to the affected cells. Systemically administered anti-restenosis drugs typically fail to achieve an effective concentration at the site of vessel injury without using unacceptably high concentrations of the drug. Delivery devices, such
10 as those described in co-owned and copending U.S. application serial no. 09/062,160, entitled "Method of Treating Restenosis by Antisense Targeting of CMV", can be used to deliver the oligomer to the site of angioplasty in a patient. The oligomer is preferably a morpholino oligomer, and is preferably contained in a pharmaceutically acceptable carrier. The oligomer may also be contained in a biocompatible polymeric carrier, also described in the above-
15 referenced application.

 Preferably, the oligomer is delivered concurrent with the angioplasty procedure. As reported by Farrell, uptake of an oligonucleotide by cells can be significantly increased when the compound is administered to balloon-injured arteries, relative to uptake by normal arteries. Drug therapy using the compositions of the invention may be combined with radiation or photodynamic
20 therapy.

 Dosages are determined according to the size of the subject, the route of administration, and the extent of the affected tissue, according to standard pharmaceutical practices. The preferred level of drug is that effective to inhibit p53 expression, and reduce or prevent restenosis, without unacceptable side effects. For an adult human, a recommended dosage is in the range of 1-25
25 μmol of antisense oligomer, and preferably 2-15 μmol . Optimum dosages for a given route can be determined by routine experimentation according to methods known in the art. For example, for delivery to the site of vessel injury, in vivo models such as described in Edelman and Rosenberg may be used.

 When the oligomer is incorporated into a drug delivery device, the device is effective to
30 deliver an appropriate dosage of drug, as described above. With respect to the surface area of tissue to be treated, an effective dose is typically in the range of 30 to 3000 μg oligomer per cm^2 of vessel wall, and more preferably about 300 to 1500 $\mu\text{g}/\text{cm}^2$. The patient may also be given the composition on a periodic basis after angioplasty, at a dosage level sufficient to further inhibit restenosis.

35

While the invention has been described with reference to specific methods and embodiments, it will be appreciated that various modifications may be made without departing from the invention.

5

EXAMPLES

Example 1

Oligonucleotide Synthesis

10 All chain-extension syntheses were performed on an Applied Biosystems Model 380B DNA synthesizer (Foster City, CA) using 1- μ mole column support and the cyanoethyl approach employing phosphoramidite chemistry as described by ABI User Bulletin, No. 58, 1991. The synthesizer was programmed with the protocol in the user's manual.

15 A summary of a typical synthesis is as follows: A 1 μ mol silica gel support column with the 3' base of the ODN (i.e. unmodified nucleotide, C-5-propyne nucleotide, or morpholino nucleotide analog) linked by the 3' hydroxy group is inserted, and synthesis is carried out in a base by base fashion from the 3' to 5' direction. All liquid reagents are supplied by Applied Biosystems, Inc. The argon gas is supplied by Air Products (Omaha). The first step is the flushing of the column with acetonitrile then argon gas which removes any residue in the column. Next, trichloroacetic acid protonates the dimethoxytrityl (DMT) group, which caps the 5' oxygen
20 of the nucleotides during the ODN synthesis. The protonated DMT is then removed from the base. The column is flushed again with acetonitrile and argon. The DMT waste is collected by the DMT waste port into test tubes and checked if it is a bright orange color. This color check is performed to insure if detritylation was successful. The DMT groups can be quantified spectrophotometrically by reading absorbency at 498 nm. If the color is not bright orange, then
25 detritylation of the bases may have not been successfully performed and synthesis of impure ODNs may occur. The next step of the synthesis is the coupling of the next base in the ODN sequence. An appropriate nucleotide base, modified to a cyanoethyl phosphoramidite, is dissolved in acetonitrile and delivered with tetrazole to the column. The bases react to form an internucleotide phosphite bond. The column is flushed with acetonitrile and argon. Any
30 uncoupled 5' hydroxy groups are capped with an acetyl group by flushing the column with acetic anhydride and 1-methylimidazole. This cap will prevent these uncoupled bases from any continuation in synthesis and reduce the length of impurities. The columns are flushed again and the trivalent phosphite is oxidized to the pentavalent phosphorothioate triester with tetraethylthiuram disulfide and acetonitrile. The column is flushed with acetonitrile and argon.
35 The synthesis is then repeated again for the next base in the sequence starting with the detritylation step. The S-ODNs are cleaved from the column by passing ammonium hydroxide

through the column and collecting the eluate. The ammonium hydroxide and cyanoethyl (from the phosphate links) are removed by evaporation of the S-ODN solution in a vacuum centrifuge for overnight. The dried ODN is diluted in sterile saline. Purity of the S-ODN is checked by running a sample of the S-ODN diluted in water on a 10% polyacrylamide gel. The
5 concentration of S-ODN is determined by reading the absorbance at 260 nm and multiplying the absorbance by its extinction coefficient.

Further details on preparation of morpholino oligonucleotide analogs can be found, for example, in Summerton and Weller, *Antisense & Nucleic Acid Drug Dev.* 7:187-195 (1997).

10

Example 2

Partial Hepatectomy

15

The partial hepatectomy study was performed on male Sprague Dawley rats (Sasco, Omaha NE) weighing between 200 - 220 grams. The animals were housed in clear plastic cages in UNMC's AAALAC- approved facility with 12 hour light/dark cycle and allowed access to Purina rat chow and tap water ad libitum. All animal protocols were approved by the 'Institutional Animal Care and Use Committee' of the university.

20

The procedure was performed as described by Higgins and Anderson (1931). Aseptic surgical techniques were employed. Rats were anesthetized using methoxyflurane (Mallinckrodt Veterinary, Mundelein, IL) and positioned with their ventral surface exposed. A three to four cm. long area was shaved along the median-line immediately posterior to the xiphoid process and swabbed with betadine. An incision was made along the median line to expose the liver. The medial and left lateral lobes were securely ligated and then excised. This resulted in removal of approximately 65 to 70 percent of the total liver. The abdominal incision was closed in two layers. Control rats that did not to receive a partial hepatectomy underwent an identical but sham
25 surgery; their livers were only exposed and not partly excised.

25

30

Oligonucleotide administration: The ODNs were injected intraperitoneally at doses of 1 mg/200gm/day S-ODN, 0.1 mg/220g/day C-5-P S-ODN, and 50nM (approx. 3.4 mg)/200g/day morpholino. All rats were administered respective ODNs immediately upon regaining consciousness after the surgery, and every 24 hr thereafter, depending on the length of the experiment. Rats were then allowed to recuperate for 1, 2, 5 or 7 days from the time of surgery. Wet weight gain was determined 24 h following administration of ODN to produce the data shown in Fig. 1.

35

Example 3

Treatment of human B-cell lymphoma.

Drug Treatment Regimen: Starting on Day 0, patients receive a morpholino antisense

oligonucleotide having SEQ ID NO:1 at a dose of 0.2 mg/kg/hr as a 7 day continuous IV infusion through a venous access device. MINE chemotherapy is given beginning on Day 4, in accordance with the following schedule:

- 5 ▪ Mesna Injection (MESNEX™, Bristol-Meyers Oncology): 500 mg/M² IV 30 minutes prior to ifosfamide, 250 mg/M² IV four hours after ifosfamide, and 250 mg/M² orally eight hours after the ifosfamide on each of days 4 - 7.
- Ifosfamide (IFEX™, Bristol-Meyers Oncology): 1.33 g/M² IV on each of days 4 - 7.
- Mitoxantrone hydrochloride (NOVANTRONE™, Lederle Laboratories) 10 mg/M² IV on day 4 only.
- 10 ▪ Etoposide (VEPESID™, Bristol-meyers Oncology) 80 mg/M² IV on each of days 4 - 7.

Patients are regularly evaluated for toxicity and response of their disease. Patients receive weekly CBC (complete blood count) and platelet counts. Cycles are repeated every 4 weeks for a maximum of 6 cycles. Patients are assessed with respect to response every cycle for disease measurable by physical examination. Patients are classified with respect to response (complete response, partial response, stable response, or progressive disease).

Patients are expected to show enhanced clinical response with this therapy as opposed to traditional MINE therapy, which does not include use of the antisense oligonucleotide SEQ ID NO:I (OL(1)p53). OL(1)p53 is expected to sensitize the cancer cells to the effects of MINE by inhibiting the cell cycle repair mechanism, thereby preferentially killing cancer cells via p53-independent apoptosis.

IT IS CLAIMED:

1. A method of treating a disease state characterized by p53 induction in a mammalian subject, comprising administering to the subject, in a suitable pharmaceutical carrier, a pharmaceutically effective amount of an antisense agent having the base sequence SEQ ID NO:1, identified as 5'-TCA GTC TGA GTC AGG CCC-3, or the base sequence SEQ ID NO:2, identified as 5'-CCC TGC TCC CCC CTG GCT CC-3',
5
7
9
11
13
15
17
19
21
23
25
27
29
31
33
35
37
39
41
43
45
47
49
51
53
55
57
59
61
63
65
67
69
71
73
75
77
79
81
83
85
87
89
91
93
95
97
99
101
103
105
107
109
111
113
115
117
119
121
123
125
127
129
131
133
135
137
139
141
143
145
147
149
151
153
155
157
159
161
163
165
167
169
171
173
175
177
179
181
183
185
187
189
191
193
195
197
199
201
203
205
207
209
211
213
215
217
219
221
223
225
227
229
231
233
235
237
239
241
243
245
247
249
251
253
255
257
259
261
263
265
267
269
271
273
275
277
279
281
283
285
287
289
291
293
295
297
299
301
303
305
307
309
311
313
315
317
319
321
323
325
327
329
331
333
335
337
339
341
343
345
347
349
351
353
355
357
359
361
363
365
367
369
371
373
375
377
379
381
383
385
387
389
391
393
395
397
399
401
403
405
407
409
411
413
415
417
419
421
423
425
427
429
431
433
435
437
439
441
443
445
447
449
451
453
455
457
459
461
463
465
467
469
471
473
475
477
479
481
483
485
487
489
491
493
495
497
499
501
503
505
507
509
511
513
515
517
519
521
523
525
527
529
531
533
535
537
539
541
543
545
547
549
551
553
555
557
559
561
563
565
567
569
571
573
575
577
579
581
583
585
587
589
591
593
595
597
599
601
603
605
607
609
611
613
615
617
619
621
623
625
627
629
631
633
635
637
639
641
643
645
647
649
651
653
655
657
659
661
663
665
667
669
671
673
675
677
679
681
683
685
687
689
691
693
695
697
699
701
703
705
707
709
711
713
715
717
719
721
723
725
727
729
731
733
735
737
739
741
743
745
747
749
751
753
755
757
759
761
763
765
767
769
771
773
775
777
779
781
783
785
787
789
791
793
795
797
799
801
803
805
807
809
811
813
815
817
819
821
823
825
827
829
831
833
835
837
839
841
843
845
847
849
851
853
855
857
859
861
863
865
867
869
871
873
875
877
879
881
883
885
887
889
891
893
895
897
899
901
903
905
907
909
911
913
915
917
919
921
923
925
927
929
931
933
935
937
939
941
943
945
947
949
951
953
955
957
959
961
963
965
967
969
971
973
975
977
979
981
983
985
987
989
991
993
995
997
999
1001
1003
1005
1007
1009
1011
1013
1015
1017
1019
1021
1023
1025
1027
1029
1031
1033
1035
1037
1039
1041
1043
1045
1047
1049
1051
1053
1055
1057
1059
1061
1063
1065
1067
1069
1071
1073
1075
1077
1079
1081
1083
1085
1087
1089
1091
1093
1095
1097
1099
1101
1103
1105
1107
1109
1111
1113
1115
1117
1119
1121
1123
1125
1127
1129
1131
1133
1135
1137
1139
1141
1143
1145
1147
1149
1151
1153
1155
1157
1159
1161
1163
1165
1167
1169
1171
1173
1175
1177
1179
1181
1183
1185
1187
1189
1191
1193
1195
1197
1199
1201
1203
1205
1207
1209
1211
1213
1215
1217
1219
1221
1223
1225
1227
1229
1231
1233
1235
1237
1239
1241
1243
1245
1247
1249
1251
1253
1255
1257
1259
1261
1263
1265
1267
1269
1271
1273
1275
1277
1279
1281
1283
1285
1287
1289
1291
1293
1295
1297
1299
1301
1303
1305
1307
1309
1311
1313
1315
1317
1319
1321
1323
1325
1327
1329
1331
1333
1335
1337
1339
1341
1343
1345
1347
1349
1351
1353
1355
1357
1359
1361
1363
1365
1367
1369
1371
1373
1375
1377
1379
1381
1383
1385
1387
1389
1391
1393
1395
1397
1399
1401
1403
1405
1407
1409
1411
1413
1415
1417
1419
1421
1423
1425
1427
1429
1431
1433
1435
1437
1439
1441
1443
1445
1447
1449
1451
1453
1455
1457
1459
1461
1463
1465
1467
1469
1471
1473
1475
1477
1479
1481
1483
1485
1487
1489
1491
1493
1495
1497
1499
1501
1503
1505
1507
1509
1511
1513
1515
1517
1519
1521
1523
1525
1527
1529
1531
1533
1535
1537
1539
1541
1543
1545
1547
1549
1551
1553
1555
1557
1559
1561
1563
1565
1567
1569
1571
1573
1575
1577
1579
1581
1583
1585
1587
1589
1591
1593
1595
1597
1599
1601
1603
1605
1607
1609
1611
1613
1615
1617
1619
1621
1623
1625
1627
1629
1631
1633
1635
1637
1639
1641
1643
1645
1647
1649
1651
1653
1655
1657
1659
1661
1663
1665
1667
1669
1671
1673
1675
1677
1679
1681
1683
1685
1687
1689
1691
1693
1695
1697
1699
1701
1703
1705
1707
1709
1711
1713
1715
1717
1719
1721
1723
1725
1727
1729
1731
1733
1735
1737
1739
1741
1743
1745
1747
1749
1751
1753
1755
1757
1759
1761
1763
1765
1767
1769
1771
1773
1775
1777
1779
1781
1783
1785
1787
1789
1791
1793
1795
1797
1799
1801
1803
1805
1807
1809
1811
1813
1815
1817
1819
1821
1823
1825
1827
1829
1831
1833
1835
1837
1839
1841
1843
1845
1847
1849
1851
1853
1855
1857
1859
1861
1863
1865
1867
1869
1871
1873
1875
1877
1879
1881
1883
1885
1887
1889
1891
1893
1895
1897
1899
1901
1903
1905
1907
1909
1911
1913
1915
1917
1919
1921
1923
1925
1927
1929
1931
1933
1935
1937
1939
1941
1943
1945
1947
1949
1951
1953
1955
1957
1959
1961
1963
1965
1967
1969
1971
1973
1975
1977
1979
1981
1983
1985
1987
1989
1991
1993
1995
1997
1999
2001
2003
2005
2007
2009
2011
2013
2015
2017
2019
2021
2023
2025
2027
2029
2031
2033
2035
2037
2039
2041
2043
2045
2047
2049
2051
2053
2055
2057
2059
2061
2063
2065
2067
2069
2071
2073
2075
2077
2079
2081
2083
2085
2087
2089
2091
2093
2095
2097
2099
2101
2103
2105
2107
2109
2111
2113
2115
2117
2119
2121
2123
2125
2127
2129
2131
2133
2135
2137
2139
2141
2143
2145
2147
2149
2151
2153
2155
2157
2159
2161
2163
2165
2167
2169
2171
2173
2175
2177
2179
2181
2183
2185
2187
2189
2191
2193
2195
2197
2199
2201
2203
2205
2207
2209
2211
2213
2215
2217
2219
2221
2223
2225
2227
2229
2231
2233
2235
2237
2239
2241
2243
2245
2247
2249
2251
2253
2255
2257
2259
2261
2263
2265
2267
2269
2271
2273
2275
2277
2279
2281
2283
2285
2287
2289
2291
2293
2295
2297
2299
2301
2303
2305
2307
2309
2311
2313
2315
2317
2319
2321
2323
2325
2327
2329
2331
2333
2335
2337
2339
2341
2343
2345
2347
2349
2351
2353
2355
2357
2359
2361
2363
2365
2367
2369
2371
2373
2375
2377
2379
2381
2383
2385
2387
2389
2391
2393
2395
2397
2399
2401
2403
2405
2407
2409
2411
2413
2415
2417
2419
2421
2423
2425
2427
2429
2431
2433
2435
2437
2439
2441
2443
2445
2447
2449
2451
2453
2455
2457
2459
2461
2463
2465
2467
2469
2471
2473
2475
2477
2479
2481
2483
2485
2487
2489
2491
2493
2495
2497
2499
2501
2503
2505
2507
2509
2511
2513
2515
2517
2519
2521
2523
2525
2527
2529
2531
2533
2535
2537
2539
2541
2543
2545
2547
2549
2551
2553
2555
2557
2559
2561
2563
2565
2567
2569
2571
2573
2575
2577
2579
2581
2583
2585
2587
2589
2591
2593
2595
2597
2599
2601
2603
2605
2607
2609
2611
2613
2615
2617
2619
2621
2623
2625
2627
2629
2631
2633
2635
2637
2639
2641
2643
2645
2647
2649
2651
2653
2655
2657
2659
2661
2663
2665
2667
2669
2671
2673
2675
2677
2679
2681
2683
2685
2687
2689
2691
2693
2695
2697
2699
2701
2703
2705
2707
2709
2711
2713
2715
2717
2719
2721
2723
2725
2727
2729
2731
2733
2735
2737
2739
2741
2743
2745
2747
2749
2751
2753
2755
2757
2759
2761
2763
2765
2767
2769
2771
2773
2775
2777
2779
2781
2783
2785
2787
2789
2791
2793
2795
2797
2799
2801
2803
2805
2807
2809
2811
2813
2815
2817
2819
2821
2823
2825
2827
2829
2831
2833
2835
2837
2839
2841
2843
2845
2847
2849
2851
2853
2855
2857
2859
2861
2863
2865
2867
2869
2871
2873
2875
2877
2879
2881
2883
2885
2887
2889
2891
2893
2895
2897
2899
2901
2903
2905
2907
2909
2911
2913
2915
2917
2919
2921
2923
2925
2927
2929
2931
2933
2935
2937
2939
2941
2943
2945
2947
2949
2951
2953
2955
2957
2959
2961
2963
2965
2967
2969
2971
2973
2975
2977
2979
2981
2983
2985
2987
2989
2991
2993
2995
2997
2999
3001
3003
3005
3007
3009
3011
3013
3015
3017
3019
3021
3023
3025
3027
3029
3031
3033
3035
3037
3039
3041
3043
3045
3047
3049
3051
3053
3055
3057
3059
3061
3063
3065
3067
3069
3071
3073
3075
3077
3079
3081
3083
3085
3087
3089
3091
3093
3095
3097
3099
3101
3103
3105
3107
3109
3111
3113
3115
3117
3119
3121
3123
3125
3127
3129
3131
3133
3135
3137
3139
3141
3143
3145
3147
3149
3151
3153
3155
3157
3159
3161
3163
3165
3167
3169
3171
3173
3175
3177
3179
3181
3183
3185
3187
3189
3191
3193
3195
3197
3199
3201
3203
3205
3207
3209
3211
3213
3215
3217
3219
3221
3223
3225
3227
3229
3231
3233
3235
3237
3239
3241
3243
3245
3247
3249
3251
3253
3255
3257
3259
3261
3263
3265
3267
3269
3271
3273
3275
3277
3279
3281
3283
3285
3287
3289
3291
3293
3295
3297
3299
3301
3303
3305
3307
3309
3311
3313
3315
3317
3319
3321
3323
3325
3327
3329
3331
3333
3335
3337
3339
3341
3343
3345
3347
3349
3351
3353
3355
3357
3359
3361
3363
3365
3367
3369
3371
3373
3375
3377
3379
3381
3383
3385
3387
3389
3391
3393
3395
3397
3399
3401
3403
3405
3407
3409
3411
3413
3415
3417
3419
3421
3423
3425
3427
3429
3431
3433
3435
3437
3439
3441
3443
3445
3447
3449
3451
3453
3455
3457
3459
3461
3463
3465
3467
3469
3471
3473
3475
3477
3479
3481
3483
3485
3487
3489
3491
3493
3495
3497
3499
3501
3503
3505
3507
3509
3511
3513
3515
3517
3519
3521
3523
3525
3527
3529
3531
3533
3535
3537
3539
3541
3543
3545
3547
3549
3551
3553
3555
3557
3559
3561
3563
3565
3567
3569
3571
3573
3575
3577
3579
3581
3583
3585
3587
3589
3591
3593
3595
3597
3599
3601
3603
3605
3607
3609
3611
3613
3615
3617
3619
3621
3623
3625
3627
3629
3631
3633
3635
3637
3639
3641
3643
3645
3647
3649
3651
3653
3655
3657
3659
3661
3663
3665
3667
3669
3671
3673
3675
3677
3679
3681
3683
3685
3687
3689
3691
3693
3695
3697
3699
3701
3703
3705
3707
3709
3711
3713
3715
3717
3719
3721
3723
3725
3727
3729
3731
3733
3735
3737
3739
3741
3743
3745
3747
3749
3751
3753
3755
3757
3759
3761
3763
3765
3767
3769
3771
3773
3775
3777
3779
3781
3783
3785
3787
3789
3791
3793
3795
3797
3799
3801
3803
3805
3807
3809
3811
3813
3815
3817
3819
3821
3823
3825
3827
3829
3831
3833
3835
3837
3839
3841
3843
3845
3847
3849
3851
3853
3855
3857
3859
3861
3863
3865
3867
3869
3871
3873
3875
3877
3879
3881
3883
3885
3887
3889
3891
3893
3895
3897
3899
3901
3903
3905
3907
3909
3911
3913
3915
3917
3919
3921
3923
3925
3927
3929
3931
3933
3935
3937
3939
3941
3943
3945
3947
3949
3951
3953
3955
3957
3959
3961
3963
3965
3967
3969
3971
3973
3975
3977
3979
3981
3983
3985
3987
3989
3991
3993
3995
3997
3999
4001
4003
4005
4007
4009
4011
4013
4015
4017
4019
4021
4023
4025
4027
4029
4031
4033
4035
4037
4039
4041
4043
4045
4047
4049
4051
4053
4055
4057
4059
4061
4063
4065
4067
4069
4071
4073
4075
4077
4079
4081
4083
4085
4087
4089
4091
4093
4095
4097
4099
4101
4103
4105
4107
4109
4111
4113
4115
4117
4119
4121
4123
4125
4127
4129
4131
4133
4135
4137
4139
4141
4143
4145
4147
4149
4151
4153
4155
4157
4159
4161
4163
4165
4167
4169
4171
4173
4175
4177
4179
4181
4183
4185
4187
4189
4191
4193
4195
4197
4199
4201
4203
4205
4207
4209
4211
4213
4215
4217
4219
4221
4223
4225
4227

13. The method of claim 12, wherein said agent is selected from the group consisting of a phosphokinase C (PKC) inhibitor, bis(chloroethyl)nitrosourea (BCNU), pentoxifylline, silymarin, staurosporine, phenylahistin, paclitaxel, retinoic acid, flavopiridol, methyl-2,5-dihydrocinnamate, herboxidiene, 9-nitrocamptothecin, maitotoxin, apigenin, nocodazole, and colcemid.

5

14. The method of claim 1, wherein said disease state results from an ischemic or ischemic/reperfusion injury.

15. A composition for use in treating a disease state characterized by p53 induction in a subject, comprising

10

an antisense agent having a base sequence selected from the group consisting of SEQ ID NO:1, identified as 5'-TCA GTC TGA GTC AGG CCC-3', and SEQ ID NO:2, identified as 5'-CCC TGC TCC CCC CTG GCT CC-3',

15

wherein the antisense agent is a morpholino oligonucleotide, a peptide nucleic acid, a 2'-O-allyl or 2'-O-alkyl modified oligonucleotide, a N3' → P5' phosphoramidate oligonucleotide, or a C-5-propyne pyrimidine-modified oligonucleotide, and a suitable pharmaceutical carrier.

20

16. The composition of claim 15, wherein the antisense agent is a morpholino oligonucleotide.

17. The composition of claim 16, wherein the morpholino oligonucleotide comprises morpholino subunits joined by phosphorodiamidate backbone linkages.

25

18. The composition of claim 15, wherein the oligonucleotide is a C-5-propyne pyrimidine-modified oligonucleotide.

19. The composition of claim 15, wherein the antisense agent has the sequence SEQ ID NO: 1.

20. The composition of claim 15, wherein the antisense agent has the sequence SEQ ID NO: 2.

30

21. An oligonucleotide having a base sequence selected from the group consisting of SEQ ID NO:1, identified as 5'-TCA GTC TGA GTC AGG CCC-3', and SEQ ID NO:2, identified as 5'-CCC TGC TCC CCC CTG GCT CC-3',

35

wherein the oligonucleotide is selected from the group consisting of a morpholino oligonucleotide, a peptide nucleic acid, a 2'-O-allyl or 2'-O-alkyl modified oligonucleotide, and a N3' → P5' phosphoramidate oligonucleotide.

22. An oligonucleotide as recited in claim 21, having the base sequence SEQ ID NO: 1.

23. An oligonucleotide as recited in claim 21, having the base sequence SEQ ID NO: 2.

24. An oligonucleotide as recited in claim 23, wherein the oligonucleotide is a morpholino oligonucleotide.

5

25. A morpholino oligonucleotide as recited in claim 24, which comprises morpholino subunits joined by phosphorodiamidate backbone linkages.

10

26. An oligonucleotide as recited in claim 23, wherein the oligonucleotide is a morpholino oligonucleotide.

27. The morpholino oligonucleotide of claim 26, which comprises morpholino subunits joined by phosphorodiamidate backbone linkages.

15

28. An oligonucleotide having a base sequence selected from the group consisting of SEQ ID NO:1, identified as 5'-TCA GTC TGA GTC AGG CCC-3', and SEQ ID NO:2, identified as 5'-CCC TGC TCC CCC CTG GCT CC-3',

wherein the oligonucleotide is a C-5-propyne pyrimidine-modified oligonucleotide.

20

29. An oligonucleotide as recited in claim 28, having the base sequence SEQ ID NO: 1.

30. An oligonucleotide as recited in claim 28, having the base sequence SEQ ID NO: 2.

1/2





Fig. 1

2/2

**Fig. 2**

SEQUENCE LISTING

<110> AVI BioPharma, Inc

<120> p53 Antisense Agent and Method

<130> 0450-0022.41

<140> Not Yet Assigned

<141> Filed Herewith

<150> US 60/105,695

<151> 1998-10-26

<160> 2

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligonucleotide

<400> 1

tcagtctgag tcaggccc

18

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligonucleotide

<400> 2

ccctgctccc ccctggctcc

20